

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
**WO 03/013597 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 39/12**,  
39/295, C12N 5/00, 15/09, 15/00, 15/74, 7/04

(21) International Application Number: PCT/US02/25185

(22) International Filing Date: 12 August 2002 (12.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/311,488 10 August 2001 (10.08.2001) US

(71) Applicant: **UNIVERSITY OF MARYLAND  
BIOTECHNOLOGY INSTITUTE** [US/US]; Office  
of Research Administration/Tech. Dev., 701 E. Pratt  
Street, Suite 200, Baltimore, MD 21202 (US).

(72) Inventor: **VAKHARIA, Vikram, N.**; 7906 Oxfarm Court,  
Bowie, MD 20715 (US).

(74) Agents: **FUIERER, Marianne** et al.; Intellectual Prop-  
erty/Technology Law, P.O. Box 14329, Research Triangle  
Park, NC 27709 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: SUB-UNIT VACCINE FOR INFECTIOUS PANCREATIC NECROSIS VIRUS

(57) Abstract: The present invention relates to sub-unit vaccines comprising structural polypeptides of Infectious Pancreatic Necrosis Virus (IPNV) comprising structural proteins V2 and V3 folded as empty IPNV viral capsid that approximates the size and structural conformation of native IPNV virus.



**WO 03/013597 A1**

## SUB-UNIT VACCINE FOR INFECTIOUS PANCREATIC NECROSIS VIRUS

### BACKGROUND OF INVENTION

#### 5 Field of the Invention

The present invention relates generally to a vaccine, and more particularly, to a sub-unit vaccine comprising structural proteins V2 and V3 of Infectious Pancreatic Necrosis Virus (IPNV) assembled as an empty viral capsid.

10

#### Description of the Related Art

Epizootics of viral infections are devastating in hatcheries and ponds rearing either cold or warm water fish and repeated disease outbreaks can jeopardize the financial survival of an operation. Thus, the health of fish is critical to the survival of the aquaculture industry and effective vaccines are desperately needed.

Infectious pancreatic necrosis virus (IPNV) is the causal agent of a highly contagious and destructive disease of juvenile Rainbow and Brook trout and Atlantic salmon. Young fish (two-to four-months old) appear to be the most susceptible to IPNV infection, resulting in high mortality. In trout and salmon, IPNV usually attacks young fry about five to six weeks after their first feeding. The affected fish are darker than usual, have slightly bulging eyes and often have swollen bellies. At the beginning of an outbreak, large numbers of slow, dark fry are seen up against water outflows, and fish are seen "shivering" near the surface. The shivering results from a characteristic symptom of the disease, a violent whirling form of swimming in which the fish rotate about their long axis. If the affected fish are examined, a characteristic white mucus is seen in the stomach. The pancreas appears to be the primary target organ for the virus.

After an IPNV outbreak, the surviving fish generally become carriers of the virus. Trout that are carriers of the virus are a serious problem for the aqua-culture industry because the only control method currently available on a commercial basis for eliminating the virus in carrier fish is destruction of these fish.

Highly virulent strains of IPNV may cause greater than 90% mortality in hatchery stocks in less than four months old. Survivors of infection can remain lifelong asymptomatic carriers and serve as reservoirs of infection, shedding virus in their feces and reproductive products. The virus is capable of infecting a number of different hosts and has a worldwide presence. IPNV can have serious economic consequences for commercial trout and salmon farms and are therefore a major concern within the aquaculture industry. Therefore, IPNV is a pathogen of major economic importance to the aquaculture industry.

10

IPNV is the prototype of the Birnaviridae virus family. IPNV contains a bisegmented dsRNA genome, which is surrounded by a single-shelled icosahedral capsid. The larger of the two genome segments, segment A (3097 bases), encodes a 106-kDa precursor polyprotein which is processed to yield mature viral structural proteins VP2 and VP3, and VP4 (also named NS) a non-structural protein (Duncan et al. 1987). VP2 has been identified as the major host protective antigen of IPNV. The genome segment B encodes a minor internal polypeptide VP1 (94 kDa) which is the putative virion-associated RNA-dependent RNA polymerase.

20 An ideal vaccine for IPNV must induce protection at an early age, prevent carrier formation, and should be effective against a large number of IPNV subtypes. One approach has been the use of killed virus as a vaccine. For example, if formalin-inactivated virus is injected intraperitoneally into four week post-hatch fry, the fish becomes immunized (Dorson, J. Virol 21:242-258, 1977). However, neither immersion of the fish into a liquid suspension of killed virus nor oral administration thereof has been found effective. Thus, the main problem with using killed virus is the lack of a practical method for administration for large numbers of immature fish because injection of the vaccine is impractical.

30 The use of attenuated viral strains have also been used as vaccines. However, the earlier attenuated strains either failed to infect the fish or failed to induce protection. Strains with low virulence have been tested as vaccines for more virulent strains, but mortality from the vaccinating strain was either too high or protection was only moderate (Hill et

al., "Studies of the Immunization of Trout Against IPN," in Fish Diseases, Third COPRAQ Session (W. Ahne, ed.), N.Y., pp. 29-36, 1980).

Recent reports have shown that expression of virus coat proteins often results in self-assembly of virus-like particles (VLP) that are essentially empty whole virions. Of these VLP-producing systems, vaccines have been proposed for poliovirus (Urakawa et al. 1989), parvovirus (Saliki et al. 1992), bluetongue virus (Belyaev et al. 1993) and infectious bursal disease virus (IBDV) - a member of the Birnaviridae family (Vakharia, et al. 1994; Bentley, et al. 1994).

10

However, several attempts have been made to recreate the same results for IPNV but to date these attempts have not been shown effective for various reasons. For instance, McKenna, et al. 2001 reported that virus like particles were generated through expression of Segment A by recombinant Semliki Forest Virus (SFV). Notwithstanding this alleged outcome, no conclusive proof was presented that the produced virus-like particles were indeed empty viral capsids. Several blots and electron microscopy slides show some type of virus like particles but without substantial proof of the formation of empty IPNV capsids resembling the size and 3D-structure of the native IPNV virus structure.

Magyar and Dobos, 1994 reported cloning of IPNV segment A into baculovirus expression vectors and expressing proteins pVP2, VP4 and VP3 in insect cells. However, as reported by Magyar and Dobos, using the baculovirus expression vectors in the insect cells did not show virus like particles that were correctly processed into a tertiary structure representing an empty viral capsid. Review of the process described in Magyar and Dobos it is clear that generating an empty IPNV capsid was impossible because Magyar and Dobos included the very first ORF of Segment A which encodes the minor 17-kDa nonstructural protein referred to as VP5 which partly overlaps the major ORF of VP2-VP-4-VP3 proteins. The VP5 protein is toxic to the cells and hence affects the production of any of the proteins. Thus, even though the proteins may have been expressed in the insect cells the proteins were not post-translationally modified and correctly folded into an empty IPNV capsid.

Phenix, et al. (2000) describes production of virus-like particles that were generated by

expressing the IPNV VP2 protein by means of a Semliki Forest Virus expression vector. However, only the VP2 protein was expressed without expressing the VP3 protein and as such, the correct formation of an empty capsid is not formed. Further, without expression of the protein VP3, aggregates may form but without the correct conformation to form neutralizing epitope. The VP2 aggregates that were formed are smaller (25nm)  
5 than virus-like particles that include a fully conformational folded viral capsid (approximately 50 to 65 nm and typically about 60nm).

Inactivated IPNV vaccines have been found to be efficacious by intraperitoneal inoculation IPNV (Leong and Fryer 1993). In addition, it was shown that the complete  
10 polyprotein of segment A expressed in *E. coli* induced protective immunity after intraperitoneal inoculation in rainbow trout fry. However, intraperitoneal inoculation for a vaccine delivery method is not very practical and bacteria are not optimal hosts for the production of many types proteins.

15 Therefore, interest has centered in other eukaryotic protein expression systems, notably yeast and insect cells in culture, as possible hosts for the production of recombinant proteins. For this reason, and related reasons, there has been effort directed toward the tissue culturing of insect cells to produce recombinant proteins. Several systems have  
20 been developed for the culture of insect cells *in vitro*, and vectors have been developed which are capable of transgene expression in insect cells. The transforming vectors are most commonly made from a group of insect pathogenic viruses belonging to the Baculoviridae family, the viruses being known as Baculoviruses. Baculoviruses are characterized by a circular double-stranded DNA genome and a rod-shaped enveloped  
25 virion. The DNA can be manipulated to incorporate a gene which encodes a subject protein and the DNA of the baculovirus will cause the cells of its host to produce the proteins encoded in its DNA.

Another approach to the production of recombinant proteins is based on the use of live  
30 insect larvae. Such an approach uses, in effect, the insect larvae as a factory for the manufacture of the desired gene product. The transgene can be expressed in the larvae through the baculovirus expression system, allowed to proliferate, and then recovered from the larvae. Because insect larvae can be grown quickly and inexpensively and the

yields obtained from insect larvae is greatly increased relative to that obtained from bacterial cells makes them an appealing alternative to cell based protein manufacturing.

Attie et al., U.S. Pat. No. 5,472,858 disclosed this approach with the tobacco hornworm.

5 After the hornworm is infected with a recombinant baculovirus, it begins secreting the recombinant protein into its hemolymph. The hemolymph can then be withdrawn using a syringe throughout the larvae's growth. However, there is a drawback to this specific method. Although the tobacco hornworm larvae is ideal for the physical manipulation because of its large size, a great deal of manual labor is required to extract the  
10 recombinant protein if large numbers are to be cultivated.

Accordingly there is a need for an IPNV sub-unit vaccine and method of producing same that overcomes the shortcomings of the prior art, that does not exhibit the problems related to live vaccine and/or attenuated vaccines, can be easily produced and recovered,  
15 and the proteins that are expressed are post-translationally modified and correctly folded into the conformation structure that exposes neutralizing epitopes.

## SUMMARY OF THE INVENTION

20

In one aspect the present invention relates to a sub-unit vaccine to reduce and/or prevent infection by Infectious Pancreatic Necrosis Virus (IPNV), the sub-unit vaccine comprising structural proteins V2 and V3 folded as an empty IPNV viral capsid. The vaccine may further comprise a reporter protein co expressed with the IPNV structural  
25 proteins.

In another aspect, the present invention relates to a baculovirus expression system comprising inclusion of a cDNA clone encoding for expressed VP2,-VP4-VP3 proteins that are self-assembled to form empty IPNV capsids that can be administered as a sub-  
30 unit vaccine.

In yet another aspect, the present invention relates to production of IPN virus-like particles having the structural conformation of native IPNV virus but without the RNA

genome.

Still another aspect of the present invention is a method of generating IPN virus-like particles assembled as an empty IPNV viral capsid, the method comprising;

5

(a) providing a recombinant baculovirus comprising a polynucleotide encoding IPNV Segment A proteins VP2,-VP4 -VP3, and a reporter protein;

(b) infecting insect larvae with the recombinant baculovirus; and

10

(c) maintaining suitable conditions for expression of IPNV Segment A proteins VP2,-VP4 -VP3, and the reporter protein to generate structural proteins VP2 and VP3 assembled as an empty IPNV capsid; and

15

(d) recovering the empty IPNV capsid from the larvae.

Another aspect relates to a sub-unit vaccine for controlling Infectious Pancreatic Necrosis Virus (IPNV) in aquatic species, the sub-unit vaccine comprising IPNV structural proteins VP2 and VP3 assembled as an empty IPNV capsid that corresponds to the 3D-  
20 structure of a native IPN virus and does not include an infectious RNA genome.

Other aspects and features of the invention will be more fully apparent from the ensuing disclosure and appended claims.

25

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is an electron microscopy slide of IPNV native full virus containing the RNA genome, the virus is sized at about 60 nm and appears white because a full virus particle, with nucleic acid, prevents stain from entering into the capsid.

30

Figure 2 is electron microscopy slide of IPNV-type particles negatively stained with uranyl acetate showing that the virus particles are empty because the stain has entered into the structure which gives a dark appearance. The virus particles have the 3D structure of native IPN viruses but show no infectious RNA genome. The particle size

corresponds to that of the native virus.

Figure 3 is a graphical representation of cumulative mortality rates of rainbow trout that were challenged with VR299

5

#### DETAILED DESCRIPTION OF THE INVENTION, AND PREFERRED EMBODIMENTS THEREOF

The present invention is based on the discovery that expression of Infectious Pancreatic  
10 Necrosis Virus (IPNV) structural proteins V2 and V3, assembled as empty viral capsids that can be administered as an effective sub-unit vaccine to reduce and/or prevent infection by IPNV.

“Sub-unit vaccine” as used herein is defined as a vaccine including sub viral components  
15 that are post-translationally modified and correctly folded to act as immunogens.

“Virus-like particles” as used herein is defined as virions that lack genetic material with 3D structure and size of a native virus.

20 “Epitopic determinants” as used herein is defined as amino acids or amino acid sequences which correspond to epitopes recognized by one or more monoclonal antibodies.

“Reporter genes” as used herein is defined as genes that express a reporter protein, which  
25 causes some determinable characteristic in a recombinant system simultaneously with the expression of the subject gene to indicate the expression of that other gene.

Expression of the Segment A gene of IPNV that encodes for VP2-VP4-VP3 by the insertion of an baculovirus expression vector leads to the production of virus-like  
30 particles formed by the self-assembly of VP2 and VP3. A cDNA clone of segment A of the IPNV consisting of a nucleotide sequence encoding for structural proteins VP2 and VP3 and a non-structural protein VP4; and a reporter gene is constructed in tandem so that the IPNV structural proteins and reporter protein are expressed simultaneously.



Briefly, the cDNA clone containing the preferred coding and/or non-coding regions of IPNV-RNA segment A can be prepared using standard cloning procedures and methods, as described for IBDV in Mundt, E., and V. N. Vakharia. 1996, Synthetic transcripts of double-stranded birnavirus genome are infectious. *Proc. Natl. Acad. Sci. USA* 93:11131-11136, the contents of which are hereby incorporated herein by reference for all purposes. Manipulations of DNAs can be performed according to standard protocols (Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning a laboratory manual*. 2nd ed. Cold Spring Harbor Laboratory. Cold Spring Harbor, N.Y.).

10

To generate cDNA clones of a coding region of the desired structural proteins, the genomic RNA is used as a template for synthesizing and amplifying according to general RT-PCR techniques well known in the art. The desired amplified fragments are then cloned into a cloning plasmid for inclusion in the baculovirus system. Preferably a reporter gene is included to simplify the harvesting and purification of the structural proteins. With this co-expression, the actual amount of subject protein produced will be directly related to the amount of reporter protein produced.

After the foreign genetic sequences for IPNV segment A and a reporter gene have been assembled, it is then necessary to package the foreign gene into a baculovirus expression vector for expression in the insect cells. This is accomplished with a transfer vector. Any commercially available Baculovirus expression kit may be used, for instance, Invitrogen Corporation markets a kit for expression of foreign genes in insect cell systems using the Baculovirus vector under the tradename MAXBAC. The kit includes Baculovirus stock, and suitable transfer vectors which may be used with the Baculovirus to transfer foreign genes into the Baculovirus for transfection into insect cells.

Baculoviruses characteristically have a circular double-stranded DNA genome which is contained in a rod-shaped enveloped virion. The DNA can be manipulated to incorporate a gene which encodes a subject protein. Like all viruses, the DNA of the baculovirus will cause the cells of its host to produce the proteins encoded in its DNA. Consequently, if the DNA of a baculovirus is manipulated to incorporate a gene which codes for IPNV VP2-VP4-VP3 protein(s) and reporter protein and that baculovirus is

allowed to infect an insect cell or insect larvae, the cells or larvae will produce the structural proteins VP2 and VP3 and the non-structural VP4.

Construction of appropriate baculovirus vectors to express a subject protein and reporter protein is apparent to one skilled in the art. The following text which is hereby incorporated herein by reference is an example of a reference that provides sufficient information and instructions to enable construction of a suitable baculovirus vector: Baculovirus Expression Vectors: A Laboratory Manual by D. R. O'Reilly, L. K. Miller and V. A. Lucklow (W. H. Freeman and Co., New York, N.Y., 1992).

10

Transfection of the baculovirus expression vector, including the gene encoding for the structural proteins of IPNV and reporter gene is transfected into cells, such as Sf9 cells, Sf21 and High Five cells, etc, which results in transcription of a recombinant baculovirus that can be used as an infectious agent to effect production of the recombinant subject and reporter proteins.

15

The recombinant baculovirus may also be used to infect insect larvae for the expression of the IPNV structural proteins and reporter proteins. Although mammalian and insect cell systems can be used to manufacture proteins, expensive and complex media are required and the bioreactors, in which the cells are grown, must be run for extended periods creating a risk of contamination of the cell culture. As such, the present invention contemplates infection of permissive insect larvae that can be infected with a baculovirus, and can be grown quickly and inexpensively. Additionally, yields can be obtained from insect larvae that cannot be obtained from cells and this fact makes insect larvae an appealing alternative to cell based protein manufacturing.

20

25

To effect infection, once a recombinant baculovirus has been constructed, a solution containing the recombinant baculovirus may be sprayed on the larvae's food for absorption therein or the baculovirus can be injected directly into the hemocoel of the larvae.

30

Because foreign proteins have been expressed in a variety of insect larvae; Bombyx mori, the silkworm (Maeda et al., 1985; Miyajima, et al., 1987), Trichoplusia ni, the cabbage

5      looper larvae (Medin et al., 1990) and *Manduca sexta*, the tobacco hornworm (U.S. Pat. No. 5,471,858) there is a wide selection of insect larvae that could be utilized with this system. The larvae of the cabbage looper has been utilized and is typical of the ideal larvae envisioned for use in the present invention. Cabbage looper larvae can be ordered from commercial sources such as Entopath, Inc. (Easton, Pa.) and can be easily grown in a laboratory according to the instructions provided by the supplier. Media for the larvae can be made from alfalfa meal, pinto beans, Brewer's yeast, ascorbic acid, wheat germ, sorbic acid, vitamins, and antibiotics (aureomycin).

10      There is a balance to be struck as to the best stage in the larval cycle of the cabbage looper or any other suitable larvae to initiate baculovirus infection. The baculovirus kills the larvae after five days so there is a limited time in which they have to grow and express the subject protein. Because the size of the larvae is related to the amount of protein they express; small larvae produce insignificant amounts of protein, it is more effective to infect the larvae when they are larger. However, if the larvae are too large, too much baculovirus is required for infection. As such, Baculovirus infection in the fourth instar, which is the last instar prior to pupation, seems to strike the optimal balance between the larvae's size and the amount of virus required for infection.

20      A preferred embodiment of the present invention is a method that infects the larvae on a continuous basis and harvesting of the larvae from the production population when the reporter gene signals production of the structural proteins. A semi-continuous embodiment is also envisioned by the inventor wherein the larvae are infected at substantially the same time and then harvested individually when they express the reporter gene.

Consequently, the reporter gene of the present invention is capable of expression in insect larvae at the same time as the gene encoding the subject protein. The determinable characteristic is a change in appearance of the living larvae that can be easily visualized.

30      Ideally, the characteristic is visible in normal light or other wavelengths of light. Thus, determination of the reporter gene's expression is simply accomplished by viewing the larvae under normal light conditions and other light conditions. Furthermore, the amount of the expressed reporter protein will directly correspond to the amount of the other

protein, such as the desired structural proteins in the larvae. Consequently, the intensity of the effect created in the appearance of the living larvae by the amount of reporter protein can be used to directly measure the amount of subject protein actually present in each larvae.

5

An excellent choice for the role of the reporter gene is the green fluorescent protein (GFP) that was originally isolated from the jellyfish, *Aequorea victoria*, and was first described in 1962 (Shimomura et al., 1962). GFP emits bright green light when simply exposed to UV or blue light, unlike other bioluminescent reporters. The emission of  
10 green light is due to the transfer of energy from the photoprotein, aequorin, of the organism to GFP. The cDNA of GFP was cloned from *Aequorea victoria* in 1992 (Prasher et al., 1992). One such GFP is a 238 amino acid protein with a molecular weight of 28 kDa (Chalfie et al., 1994) having a major absorption peak at 395 nm and a minor peak at 470 nm with a single emission peak at 509 nm (Chalfie et al., 1994).  
15 Preferably, the GFP gene has a nucleotide sequence selected from the group consisting of SEQ ID.NOs: 1, 8, or 9.

Advantageously, fluorescence of GFP is species-independent and requires no substrate, cofactor, or additional proteins for illuminating green light. Unlike other reporter tags  
20 such as luciferase, b-galactosidase, or fluorescent-tagged antibodies, GFP does not require fixation techniques that are toxic to the cells under the investigation (Chalfie et al., 1994).

Additionally, GFP mutations have also been developed that serve well in the capacity of  
25 the reporter gene. GFPuv was optimized for UV excitation (Cramer et al., 1996). GFPuv is 18 times brighter than wild-type GFP and can be easily detected by the naked eye when excited with standard, long-wave UV light (e.g., source for many DNA transilluminator light tables). This variant contains additional amino acid mutations which increases its translational efficiency. Purified GFPuv has the same excitation and  
30 emission maxima as wild-type GFP.

Mutants of GFP are available commercially and include variants with the blue and red-shifted proteins along with several that have various intensities of green for which the

codon composition has been altered . Among the commercial vendors of these mutants are Life Technologies, Inc., Clontech, Inc., and Invitrogen, Inc.

It is envisioned that the present invention will include the use of an automated system for  
5 selecting individual larvae for harvest. A monitoring capability could be added to such a system by attaching an LED/detector to each fin where the larvae hang and wiring the LED/detector and a fin to a separate controller. When the LED/detector detects a sufficient intensity of the green color of the GFP expressed in a larva, a signal would be sent to the controller which in turn would deliver voltage to the fin where the  
10 transmitting LED/detector is located. The voltage would shock the larva causing it to fall off the fin into a collection receptacle. Another possible monitoring system could resemble the conveyor belt/switching gate apparatus used by dairy egg manufacturers, where the brightest larvae are harvested.

15 Although an automated system is preferable, since GFP mutants expressed in larvae, such as GFPuv, can be discerned by the human eye in normal laboratory light in larvae at least two or three days after infection with baculovirus and detection of GFP with human eyesight under UV light possible after at least three days, manual infection and selection for harvest is certainly feasible.

20

Conditions which enhance the expression of proteins VP2, VP4 and VP3 and the reporter gene, particularly green fluorescent protein, include infection with a viral loading of at least  $5 \times 10^7$  pfu/mL recombinant baculovirus, temperature of at least about 30°C, harvesting of the larvae at least 3-5 days, more particularly at least 4 days, after post  
25 infection; at a pH of at least about 3.5 to about 4.0 and in the presence of protease inhibitors such as PMSF, EDTA and benzamidine, preferably at least about 1.5 mM.

Preservation of the larvae after selection for harvest in the present invention is envisioned to be conducted by freezing the selected larvae and then homogenizing the larvae in  
30 conditions which minimize the activity of proteases.

The previously described versions of the present invention have many advantages including the easy selection of larvae -at the point of their optimal protein expression.

Because the visible expression of GFP will occur simultaneously with the expression of the IPNV structural proteins, each larvae can be selected for harvest when it is expressing the optimal amount of the subject protein. Furthermore, if the GFP is fused with at least one of the structural proteins, when expressed, it is possible in the present invention to  
5 quantify the amount of subject protein in an individual larvae. This in turn allows an estimation of the total yield from a production population to be made.

In addition, larvae which are not expressing significant amounts of the IPNV structural proteins can be removed from the production population so that resources are not  
10 expended attempting to purify the subject protein from them. Furthermore, because the larvae can be grown on inexpensive media, it is unnecessary to incur the high expense of formulating complex media and maintaining bioreactors.

The visualization of the reporter protein will permit the progress of the expression of V2-  
15 V4-V3 polypeptide to be monitored through the purification process directly and indirectly. This will also serve to facilitate purification. Separation of the reporter protein from the V2-V4-V3 polypeptide may be accomplished in several ways. For example, an affinity ligand could be engineered onto the reporter protein. The affinity ligand can be used to bind and separate the structural proteins from the larval extract  
20 during the initial purification.

As a further means to simplify purification, the linkage between the gene for the V2-V4-  
V3 polypeptide and the reporter gene could comprise a gene which expresses a protein that is cleaved by a specific enzyme. Once the fusion product is separated from the  
25 homogenate via the affinity ligand, it could be exposed to the enzyme which cleaves the linking protein to separate the subject protein and the reporter protein. Then, a one step purification could be performed to purify the subject protein.

Still further, in the present invention with the inclusion of the VP4 protease as part of the  
30 subject protein, it is contemplated that the VP4 may participate in the release of the GFP protein from either of the V2 or V3 structural proteins.

Once the V2-V4-V3 polypeptide and/or V2 and V3 proteins are separated from the

reporter protein, if it is determined that the GFP protein must be removed for effective vaccine, the purified VLPs are administered as a sub-unit vaccine to aquatic species.

5 It is contemplated by the inventors to include more than one strain of IPNV so that more than one type of empty viral capsid can be generated and recovered. Thus, the recovered empty viral capsids may contain epitopic determinants for more than one strain of IPNV. Since VP2 protein is the major host protective immunogen of IPNV, the present invention can include structural proteins (VP2 and VP3) from at least two different IPNV strains. Strains suitable for use in producing the present vaccine include but are not  
10 limited to West Buxton, Jasper, SP, N1, DRT, Ab, HE, TE, Canada 1, 2, 3 and VR299 strains.

Physiologically acceptable carriers for vaccination of fish are known in the art and need not be further described herein. In addition to being physiologically acceptable to the  
15 fish the carrier must not interfere with the immunological response elicited by the vaccine and/or with the expression of its polypeptide product.

Other additives, such as adjuvants and stabilizers, among others, may also be contained in the vaccine in amounts known in the art. Preferably, adjuvants such as aluminum  
20 hydroxide, aluminum phosphate, plant and animal oils, and the like, are administered with the vaccine in amounts sufficient to enhance the immune response to the IPNV. The amount of adjuvant added to the vaccine will vary depending on the nature of the adjuvant, generally ranging from about 0.1 to about 100 times the weight of the IPNV, preferably from about 1 to about 10 times the weight of the IPNV.

25 The vaccine of the present invention may also contain various stabilizers. Any suitable stabilizer can be used including carbohydrates such as sorbitol, mannitol, starch, sucrose, dextrin, or glucose; proteins such as albumin or casein; and buffers such as alkaline metal phosphate and the like.

30 The vaccine can be administered by any suitable known method of inoculating fish including but not limited to immersion, oral administration, spraying and injection. Preferably, the vaccine is administered by mass administration techniques such as

immersion as conducted by a standardized immersion protocol described by McAllister and Owens (1986), the contents of which are hereby incorporated by reference herein in its entirety.

- 5 When administered by injection, the vaccines are preferably administered parenterally. Parenteral administration as used herein means administration by intravenous, subcutaneous, intramuscular, or intraperitoneal injection. Further administration may be accomplished by sonification or electroporation.
- 10 The vaccine of the present invention is administered to fish to prevent IPNV anytime before or after hatching. The term "fish" is defined to include but not be limited to fish species including trout, salmon, carp, perch, pike, eels, and char as well as mollusks and crustaceans.
- 15 The vaccine may be provided in a sterile container in unit form or in other amounts. It is preferably stored frozen, below  $-20^{\circ}\text{C}$ , and more preferably below  $-70^{\circ}\text{C}$ . It is thawed prior to use, and may be refrozen immediately thereafter. For administration to fish, the recombinantly produced VLPs may be suspended in a carrier in an amount of about  $10^2$  to  $10^7$  pfu/ml, and more preferably about  $10^5$  to  $10^6$  pfu/ml in a carrier such as a saline
- 20 solution. The sub-unit vaccine may contain the antigenic equivalent of  $10^4$  to  $10^7$  pfu/ml suspended in a carrier. Other carriers may also be utilized as is known in the art.

Examples of pharmaceutically acceptable carriers are diluents and inert pharmaceutical carriers known in the art. Preferably, the carrier or diluent is one compatible with the

25 administration of the vaccine by mass administration techniques. However, the carrier or diluent may also be compatible with other administration methods such as injection, and the like.

The invention also can be used to produce combination vaccines wherein the IPNV

30 material is combined with antigen material of other relevant fish pathogens and/or bacterial antigens. Examples of relevant fish pathogens include but are not limited to infectious hematopoietic necrosis virus (IHNV), viral hemorrhagic septicemia virus (VHSV), ISAV (Infectious salmon anemia virus), PDV (Pancreas disease virus), Irido



virus and Nodavirus. Examples of relevant bacterial antigens include but are not limited to antigens from gram positive bacteria such as but not limited to *Lactococcus garvieae* and gram negative bacteria such as but not limited to *Aeromonas salmonicida*. Other relevant bacterial antigens include but are not limited to antigens from *Vibrio anguillarum*, *Vibrio salmonicida*, *Vibrio viscosus*, *Yersinia ruckri*, *Piscirickettsia salmonis*, *Renibacterium salmoninarum*, *Pasteurella piscicida*, *Flavobacterium columnare*, and *Flavobacterium psychrophilum*.

The foregoing embodiments of the present invention are further described in the following Examples. However, the present invention is not limited by the Examples, and variations will be apparent to those skilled in the art without departing from the scope of the present invention.

**Cloning and expression of ALV122 segment A (major ORF) and EGFP in Bac-to-Bac baculovirus expression system.**

All DNA manipulations were carried out according to standard molecular biology techniques described by Sambrook, et al. A full length complementary DNA fragment (SEQ ID NO: 2) encoding the Segment A (V2-V4-V3) of a Norwegian field isolate (Sp serotype (ALV122)) of Infectious Pancreatic Necrosis Virus (IPNV) was generated by reverse transcription-polymerase chain reaction (RT-PCR).

To generate cDNA clones of segment A of Sp strain (ALV122) (SEQ ID NO: 2), two primer pairs (A-A5' NC plus SpA-KpnR, SpA-KpnF plus SpA-PstR) were used for RT-PCR amplification. The sequences of these primers were:

- 1) A-A5' NC, 5'-TAATACGACTCACTATAGGAAAGAGAGTTTCAACG-3' (SEQ ID NO: 10);
- 2) SpA-KpnR, 5'-GGCCATGGAGTGGTACCTTC-3' (SEQ ID NO: 11);
- 3) SpA-KpnF, 5'-GAAGGTACCACTCCATGGCC-3' (SEQ ID NO: 12; and
- 4) SpA-PstR, 5'-AAAGCTTCTGCAGGGGGCCCCCTGGGGGGC-3' (SEQ ID NO: 13).

Using genomic RNA as a template, desired overlapping cDNA fragments of segment A were synthesized and amplified according to the supplier's protocol (Perkins-Elmer). Amplified fragments were cloned into the EcoRI site of pCR2.1 vector (Invitrogen Corp.) to obtain plasmids pCRSpA5' and SpA31. The insert DNA in all the plasmids  
5 was sequenced by the dideoxy chain termination method using an Applied Biosystem automated DNA sequencer, and the sequence data was analyzed by using PC/GENE (Intelligenetics) software. To construct a full-length cDNA clone of segment A, a representative plasmid of pCRSpA5' and pCRSpA3' clones was double-digested with restriction enzyme pairs BamHI plus KpnI and KpnI plus HindIII release 1495 and 1602  
10 bp fragments, respectively. These fragments were then cloned between the BamHI and KpnI sites of pUC19 vector to obtain plasmid PUC19SpAALV122#7. This plasmid contained a full-length copy of segment A which encodes all for VP2-VP4-VP3.

A complementary DNA clone of energetic GFP (SEQ ID NO: 1) was amplified using  
15 primers XhoEGFPF: 5'-AACTCGAGATGGTGAGCAAGGGCGAG-3' (SEQ ID NO: 4) and XhoEGFPR: 5'-ATCTCGACTTGTACAGCTCGTCCATGC-3' (SEQ ID NO: 5). The PCR product was cloned into pCR 2.1 vector by TA cloning. (A TOPO TA cloning kit is available from InVitrogen Corp containing T vector and other components required for cloning including the pCR2.1-TOPO vector, 10X PCR buffer, salt solution, dNTP  
20 mix, control template, and control PCR primers, DH5a-T1 Competent cells (1 vial/transformation), SOC medium.) EGFP was excised using Xho I site and cloned into a pFastBac DUAL vector next to the P10 promoter to yield FastBacEGFP(p10).

The construction of the full-length cDNA clone of segment A of IPNV strain ALV103 of  
25 Sp serotype has been described in U.S. Patent No. 6,274,147, the contents of which are hereby incorporated herein by reference for all purposes. Using the methods described in U. S. Patent No: 6,274,147, the major open reading frame (ORF) of segment A was amplified by PCR with the following set of primers.: SpABamF: 5'-GGGATCCATGAACACAAACAAGGC-3' (SEQ ID NO: 6) and SpAHinR: 5'-  
30 AAAGCTTACACCTCAGCGTTGTC-3' (SED ID NO: 7). The PCR product was cloned into pCR2.1 vector by TA cloning.

The SP strain ALV103 was cloned behind the polyhedrin promoter between the BamHI and HindIII sites of baculovirus vector, pBlueBac4. The recombinant plasmid BlueBacSPA#8 was obtained. This plasmid was digested with BstE II and Hind III enzymes and this fragment was replaced with a BstE II and Hind III fragment from plasmid pUC19ALV122A #7 containing the entire segment A of ALV122. The resulting plasmid carrying the major ORF of segment A from ALV122 strain was digested with BamHI and Sal I enzymes and cloned next to polyhedrin promoter of FastBacEGFP(p10) to yield FastBacEGFP(p10)IPNA(poly). This plasmid was then used to make bacmid clone and subsequently used to generate recombinant baculovirus.

10

A recombinant baculovirus containing the nucleotide sequence of segment A of IPNV and the EGFP gene was obtained by the method described in the manual of O'Reilly et al. (1991) and using a BAC-TO-BAC Baculovirus Expression system available from Invitrogen. The system includes pFASTBAC™ the BAC-TO-BAC expression vector for transforming DH10Bac E.coli which contains a specialized Bacmid that recombines with the preferred construct via site-specific transposition to create a recombinant expression Bacmid.

The mixture was added to 0.75 ml of Grace's medium supplemented with 10% FBS in a 60-mm dish seeded with the permissive *Spodoptera frugiperda* (Sf9) cells. Following incubation at 27°C for 4 hr, the medium was removed; the monolayer washed with Grace's medium supplemented with 10% FBS and the dish incubated at 27°C. Four to six days post transfection, the cells were observed with an inverted microscope for signs of infection. Extracellular virus was collected and plaqued on monolayer of Sf9 cells. Representative recombinant IPNV was used for infecting of insect larvae.

**Infecting Insect Larvae to generate sufficient quantities of empty IPNV capsids for vaccination purposes.**

This example describes the optimized production of IPNV structural proteins in larvae of the cabbage looper, *Trichoplusia ni*. The eggs were obtained from a commercial supplier (Entopath, Inc., Easton, Pa.) and hatched in Styrofoam cups containing solid food (Entopath) at 30°C. The recombinant baculovirus, isolated and purified from the insect

cells were used to infect the larvae by injection of 5  $\mu$ l of  $5 \times 10^6$  pfu/ml per fish. (It should be noted that infection may also be accomplished by spreading 500  $\mu$ L recombinant baculovirus ( $5 \times 10^7$  pfu/ML) on the media which was obtained pre-made in Styrofoam cups (Entopath)).

5

The cups were covered and allowed to stand an hour for the virus to be completely absorbed by the media. The fourth instar larvae (about 4 days after hatching) were then placed into the cups (approximately about 10 to 15 larvae per cup). The cups were then inverted and the larvae were allowed to feed on the infected food at 30°C. The fecal matter dropped onto the lid so it could be discarded daily.

10

The infected larvae were then collected and frozen at about -60°C until they were ready for isolation and purification of the recovered protein structures. The frozen larvae were thawed, and homogenized in phosphate buffered saline (PBS) containing 60 mM dithiothreitol (DTT), and 0.5% Triton X-100 at pH 7.0. The homogenate was then centrifuged at 4°C. to remove large debris. After centrifugation, the supernatant was also further clarified with using a 0.22 micron filter.

15

The fraction containing the expressed structural proteins was examined in an electron microscope. Empty IPNV capsids were found as shown in Figure 2, that were similar to size and symmetry to those described for native ALV122 Sp strains as shown in Figure 1. The particles viewed under the electron microscope were sized at about 60 nm and showing the capsid structure but having no RNA genome as compared to the fully active native IPN viruses. Thus, the shape and size of the negatively stained purified IBNV-like particles were similar to in size and shape to the native IPNV but without the infectious genomic load.

25

The virus particles were recognized by anti-VP2 monoclonal antibody provided in an antibody kit commercially available from Dioxotics, which further confirmed the identity with the native IPNV and suggested that the surface of the virus particles were formed by VP2 proteins. Further binding of the antibody to the virus particles were evidence of correct structural formation of the outer viral capsid which has internalized the VP3 protein. As such, the VP2 protein, carrying the major neutralizing epitope formed the

30

external surface of the virion and the VP3 protein faces the interior of the capsid.

#### **Efficacy of empty IPNV capsids as a sub-unit vaccine**

5 This study was carried out at the United States Geological Survey, National Fish Health  
Research Laboratory, Kearneysville, West Virginia. The experiment was conducted in a  
flow through system. The experimental design consisted of four experiments with six  
treatments and 3 replicates with a total of seventy-two tanks. Thirty Rainbow trout fry,  
with an average body weight of 0.13 g were held in one-liter polypropylene tanks and  
10 used for the study. In each experiment, fish was vaccinated at high (0.5mg/mL) and low  
(0.1mg/mL) doses of the expressed IPNV structural proteins in three treatments and the  
other three treatments was treated with phosphate buffered saline (PBS) instead, and  
served as controls. In the first and second experiment, fish were subjected to high dose  
of the IPNV structural proteins and were challenged after two and three weeks post  
15 vaccination respectively, with Sp and VR-299 strains of IPNV. Low dose of expressed  
structural IPNV protein was used in the third and fourth experiment with similar post  
vaccination exposure times. Overall, the experimental set up consisted of 72 tanks (6  
groups X 3 replicates X 2 vaccine doses X 2 time intervals = 72). The treatment groups  
and the number of tanks that were used for the study are shown in Table 1.

20

Table 1. Treatment groups and the total number of tanks used for the study. Control  
groups were treated with phosphate buffered saline (PBS) instead of expressed IPNV  
structural proteins.

Treatment	Replicates	Vaccine Dose	Exposure time	Total Tanks
<b>Control Groups</b>				
No vaccine + No Challenge	3	2 (PBS)	2	12
No Vaccine + ALV122 (Sp)	3	2 (PBS)	2	12
No Vaccine + VR-299	3	2 (PBS)	2	12
<b>Immunized Groups</b>				
Immunized + No challenge	3	2	2	12
Immunized + ALV122	3	2	2	12

Immunized + VR-299	3	2	2	12
--------------------	---	---	---	----

For vaccination and challenging of the fish, a standardized immersion challenge described by McAllister and Owens (1986) for IPNV was followed. For vaccination, the purified larval homogenate containing expressed IPNV proteins was used. For virus challenge, stock virus was diluted in PBS and added to tanks containing fish at a density of 1g of fish per 25 mL of water to achieve a concentration of  $10^5$  PFU/mL. During both vaccination and challenge, fish were exposed for 5 hours with static condition and aeration. Water flow was resumed at a rate of 250 mL/min after the end of exposure.

Mortality was monitored over a period of 28 days in all the four experiments. The dead fish were collected daily and frozen at  $-20^{\circ}$  C until the analysis. At the end of all the experiments, seventy-five fish from both control and vaccinated groups (twenty-five from each replicate) including the survivors and the dead fish were individually analyzed for the presence of IPNV by viral plaque assay. Histopathological studies also will be performed to examine whether the vaccine can prevent lesions in the internal organs.

## Results

In experiment 1, fish were immunized with high dose of IPNV expressed structural proteins and challenged two weeks post-vaccination with Sp and VR299 strains of IPNV. Mortality due to primary infection started to appear on the seventh day in the control group that was not immunized but challenged with VR299. Mortality in the immunized group on the other hand, started appearing on the thirteenth day in the VR299 challenged group, which indicates that the vaccine was able to prevent the primary infection. It was found that the cumulative mortality rate in the control group was 13.5% while in the immunized group it was only 8 % (Fig 3). The result indicated that the expressed proteins of Sp strain that was isolated in Norway can confer a partial cross protection against VR299 strain that was isolated from a field outbreak in USA. Neither the control nor the treated group showed mortality when exposed to the Sp strain of IPNV. The Sp isolate that was used in the study was originally isolated from a field outbreak in Norway from Atlantic salmon. The results obtained indicate that the rainbow trout used in the study might not have the receptors for the virus entry and hence may not be susceptible to Sp strain.

The lower doses used in Experiments 3 and 4 showed no difference in the mortality rate indicating that the effectiveness of the vaccine is dose dependent.

## REFERENCES

All cited references are hereby incorporated herein by reference for all purposes.

- 5 Belyaev, A.S. and P. Roy. 1993 Development of baculovirus triple and quadruple expression vectors: co-expression of three or four bluetongue virus proteins and the synthesis of bluetongue virus-like particles in insect cells. Nucl. Acids. Res. 21(5):1219-1223.
- 10 Bentley, WE., M.Y. Wang and V.N. Vakharia. 1994. Development of an efficient bioprocess for poultry vaccines using high-density insect cell culture. Ann. New York Acad. Sci. 745: 336-359.
- 15 Cha, HJ., N.G. Dalal, M.Q. Pham, V. N. Vakharia, G. Rao, WE. Bentley. 1999. Insect larval expression process is optimized by generating fusions with green fluorescent protein. Biotechnol. Bioeng. 65:316-324.
- 20 Duncan, R., E. Nagy, P.J. Krell and P. Dobos. 1987. Synthesis of the infectious pancreatic necrosis virus polyprotein, detection of a virus-encoded protease, and line structure mapping of genome segment A coding regions. 3. Virol. 61:3655-3664.
- 25 Leong, J.C. and J.L. Fryer. 1993. Viral vaccines for aquaculture. Pages 225-240 in M. Faisal and F.M. Hetrick, eds. Annual Review of Fish Diseases. Vol.3. Pergamon Press, New York.
- Maeda, S. 1989. Expression of foreign genes in insects using baculovirus vectors. Ann. Rev. Entomol. 34:351-72.
- 30 Magyar, G., and Dobos, P. (1994). Expression of infectious pancreatic necrosis virus polyprotein and VP1 in insect cells and the detection of the polyprotein in purified virus. Virology 198: 437-445.



- McAllister P.E., Owens, W.J., Infectious Pancreatic Necrosis Virus - Protocol for a Standard Challenge to Brook Trout, Transactions of the American Fisheries Society, 115 (3): 466-470 May 1986.
- 5 McKenna, B.M., Fitzpatrick, R.M., Phenix, K.V., Todd, D., Vaughan, L.M., and Atkins, G.J., 2001. Formation of Infectious Pancreatic Necrosis Virus-like Particles Following Expression of Segment A by Recombinant Semliki Forest Virus, Mar. Biotechnol. 3, 103-110.
- 10 O'Reilly, D.K., L.K. Miller and VA Luckow. 1991. Baculovirus expression vectors: A laboratory manual. 1<sup>st</sup> edition. W.H. Freeman and Co., New York.
- Phenix, K.V., McKenna, B., Fitzpatrick, R., Vaughan, L., Atkins, G., Liljestrom, P and Todd, D., 2000. Cell Culture Evaluation of the Semliki Forest Virus Expression System  
15 As a Novel Approach for Antigen Delivery and Expression in Fish, Mar. Biotechnol. 2, 27-37.
- Saliki, J.T., B. Mizak and H.P. Flore. 1992. Canine parvovirus empty capsids produced by expression in a baculovirus vector: use in analysis of viral properties and  
20 immunization. J. Gen. Virol. 73:369.
- Sambrook, J., E.F. Fritsch and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2<sup>nd</sup> ed Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.  
25
- Urakawa, T., M. Ferguson, P.D. Minor, J. Cooper, M. Sullivan, J.W. Almond and D.H.L. Bishop. 1989. Synthesis of immunogenic, but non-infectious, poliovirus particles in insect cells by a baculovirus expression vector. J. Gen. Virol. 70:1453-1463.
- 30 Vakharia, V.N., D.B. Snyder, D. Luticken, S.A. Mengel-Whereat, P. K. Savage, G. H. Edwards and MA Goodwin. 1994. Active and passive protection against variant and classic infectious bursal disease virus induced by baculovirus expressed structural proteins. Vaccine 12:452-456.

Yao, K. and V.N. Vakharia 1998. Generation of infectious pancreatic necrosis virus from cloned cDNA J. Virol. 72:8913-8920.

5

## THE CLAIMS

What is claimed is:

- 5 1. A sub-unit vaccine for controlling Infectious Pancreatic Necrosis Virus (IPNV) in aquatic species, the sub-unit vaccine comprising IPNV structural proteins VP2 and VP3 assembled as an empty IPNV capsid.
2. The vaccine according to claim 1, further comprising a green fluorescent protein.
- 10 3. The vaccine according to claim 1, wherein the IPNV is a strain selected from the group consisting of West Buxton, Jasper, SP, N1, DRT, Ab, HE, TE, Canada 1, Canada 2, Canada 3 and VR299 strains.
- 15 4. The vaccine according to claim 3, further comprising an antigen from an aquatic virus other than IPNV selected from the group consisting of: infectious hematopoietic necrosis virus (IHNV), viral hemorrhagic septicemia virus (VHSV), ISAV (Infectious salmon anemia virus), PDV (Pancreas disease virus), Irido virus, and Nodavirus.
- 20 5. The vaccine according to claim 1, wherein the empty IPNV capsid approximates the size and conformation of a native IPNV virus.
6. The vaccine according to claim 1, wherein the empty viral capsid has a diameter of about 50 to about 65 nm.
- 25 7. The vaccine according to claim 1, wherein the VP2 and VP3 structural proteins are encoded by SEQ ID NO: 2.
8. The vaccine according to claim 1, further comprising a physiologically acceptable
- 30 carriers for fish.
9. A baculovirus expression vector comprising a polynucleotide sequence encoding for structural proteins VP2-VP4-VP3 of infectious pancreatic necrosis virus and a green

fluorescent protein.

10. The baculovirus expression vector according to claim 9, wherein the polynucleotide sequence encoding for the infectious pancreatic necrosis virus is SEQ ID NO: 2, and the green fluorescent protein is SEQ ID NO: 1.

11. The baculovirus expression vector according to claim 9, wherein the polynucleotide sequence encoding for the green fluorescent protein is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 8, and SEQ ID NO: 9.

10

12. A host cell transfected with the baculovirus expression vector according to claim 9.

13. A host cell transfected with the baculovirus expression vector according to claim 10.

14. A host cell transfected with the expression vector according to claim 11.

15. The host cell according to claim 10, wherein the host cell is an insect cell.

20

16. A method of generating structural proteins of IPNV assembled as an empty viral capsid comprising the steps of:

(a) providing a recombinant baculovirus comprising a polynucleotide encoding IPNV Segment A proteins VP2,-VP4 -VP3, and a reporter protein;

(b) infecting insect larvae with the recombinant baculovirus; and

(c) maintaining suitable conditions for expression of IPNV Segment A proteins VP2,-VP4-VP3, and the reporter protein to generate structural proteins VP2 and VP3 assembled as an empty IPNV capsid; and


(d) recovering the empty IPNV capsid from the larvae.

- 17: The method according to claim 16, wherein the larvae are infected all at the same time and harvested when the reporter protein is expressed.
- 5 18. The method according to claim 16, wherein suitable conditions comprises:  
infesting the larvae with a viral loading of at least  $5 \times 10^7$  pfu/mL recombinant baculovirus;  
maintaining a temperature of at least about 30°C; and  
harvesting of the larvae at least 3-5 days after post infection at a pH of at least about 3.5  
10 to about 4.0.
19. A method for reducing and/or preventing infection of IPNV in marine fish by administering an effective amount of a sub-unit vaccine comprising IPNV structural proteins VP2 and VP3 assembled as an empty IPNV capsid that approximates the size  
15 and conformation of a native IPN virus.
20. The method according to claim 19, wherein the IPNV is a strain selected from the group consisting of West Buxton, Jasper, SP, N1, DRT, Ab, HE, TE, Canada 1, Canada 2, Canada 3 and VR299 strains.
- 20 21. The method according to claim 19, wherein the vaccine further comprising an antigen from an aquatic virus other than IPNV selected from the group consisting of: infectious hematopoietic necrosis virus (IHNV), viral hemorrhagic septicemia virus (VHSV), ISAV (Infectious salmon anemia virus), PDV (Pancreas disease virus), Irido  
25 virus, and Nodavirus.
22. The method according to claim 19, wherein the empty viral capsid resembles the 3D-structure of native IPNV particles and does not include an infectious RNA genome.
- 30 23. The method according to claim 19, wherein the empty IPNV capsid has a diameter of about 50 to about 65 nm.
24. A sub-unit vaccine for controlling Infectious Pancreatic Necrosis Virus (IPNV) in

aquatic species, the sub-unit vaccine comprising IPNV structural proteins VP2 and VP3 assembled as an empty IPNV capsid that corresponds to the 3D-structure of a native IPN virus and does not include an infectious RNA genome.


5

1/3



**Figure 1**

2/3



**Figure 2**






Figure 3

## SEQUENCE LISTING

&lt;110&gt; VIKHARIA, VIKRAM N

&lt;120&gt; SUB-UNIT VACCINE FOR INFECTIOUS PANCREATIC NECROSIS VIRUS

&lt;130&gt; 4115-154

&lt;150&gt; 60/311,488

&lt;151&gt; 2001-08-10

&lt;160&gt; 13

&lt;170&gt; PatentIn version 3.1

&lt;210&gt; 1

&lt;211&gt; 720

&lt;212&gt; DNA

&lt;213&gt; Aequorea victoria

&lt;400&gt; 1

```

atggtgagca agggcgagga gctgttcacc ggggtggtgc ccatcctggt cgagctggac      60
ggcgacgtaa acggccacaa gttcagcgtg tccggcgagg gcgagggcga tgccacctac      120
ggcaagctga ccctgaagtt catctgcacc accggcaagc tgcccgtgcc ctggcccacc      180
ctcgtgacca ccctgacctc cggcgtgcag tgcttcagcc gctaccccga ccacatgaag      240
cagcacgact tcttcaagtc cgccatgccc gaaggctacg tccaggagcg caccatcttc      300
ttcaaggacg acggcaacta caagaccgcg gccgaggtga agttcgaggg cgacaccctg      360
gtgaaccgca tcgagctgaa gggcatcgac ttcaaggagg acggcaacat cctggggcac      420
aagctggagt acaactacaa cagccacaac gtctatatca tggccgacaa gcagaagaac      480
ggcatcaagg tgaacttcaa gatccgccac aacatcgagg acggcagcgt gcagctcgcc      540
gaccactacc agcagaacac ccccatcggc gacggccccg tgctgctgcc cgacaaccac      600
tacctgagca ccagtcgcgc cctgagcaaa gaccccaacg agaagcgcga tcacatggtc      660
ctgctggagt tcgtgaccgc cgccgggatc actctcgga tggacgagct gtacaagtaa      720

```

&lt;210&gt; 2

&lt;211&gt; 3097

&lt;212&gt; DNA

&lt;213&gt; Infectious pancreatic necrosis virus

&lt;400&gt; 2

```

ggaaagagag tttcaacggt agtggttaacc cagagcgga gagctcttac ggaggagctc      60
tccgtcgatg gcgaaagccc tttctaacaa aaaaacaaac aatctatatc aatgcaagat      120
gaacacaaac aaggcaaccg caacttacct gaaatccatt atgcttccag agactggacc      180
agcaagcatc ccggacgaca taacggagag acacatctta aaacaagaga cctcgtcata      240
caacttagag gtctccgaat caggaagtgg cattcttgtt tgtttccctg gggcaccagg      300
ctcacggatc ggtgcacact acagatggaa tgcaaccag acggggctgg agttcgacca      360
gtggctggag acgtcgcagg acctgaagaa agccttcaac tacgggaggc tgatctcaag      420
gaaatatgac atccaaagct ccacactacc ggccgggtctc tatgctctga acgggacgct      480
caacgctgcc accttcgaag gcagtctgtc tgagggtggag agcctgacct acaacagcct      540

```

gatgtcccta acaacgaacc cccaggacaa agtcaacaac cagctggtga ccaaaggagt	600
cacagtccctg aatctaccaa cagggttcga caaaccatac gtccgcctag aggacgagac	660
accccagggt ctccagtcaa tgaacggggc caagatgagg tgcacagctg caactgcacc	720
gcgagggtac gagatcgacc tcccatccca acgcctaccc cccgttactg cgacaggaac	780
cctcaccact ctctacgagg gaaacgccga catcgtcaac tccacgacag tgacgggaga	840
cataaacttc agtctgacag aacaaccgcg agtcgagacc aagttcgact tccagctgga	900
cttcatgggc cttgacaacg acgtcccagt tgtcacagtg gtcagctccg tgctggccac	960
aaatgacaac tacagaggag tctcagccaa gatgaccagc tccatcccga ccgagaacat	1020
cacaaagccg atcaccaggg tcaagctgtc atacaagatc aaccagcaga cagcaatcgg	1080
caacgtcgcc accctgggca caatgggtcc agcatccgtc tccttctcat cagggaacgg	1140
aaatgtcccc ggcgtgctca gaccaatcac actggtggcc tatgagaaga tgacaccgct	1200
gtccatcctg accgtagctg gagtgtccaa ctacgagctg atcccaaacc cagaactcct	1260
aaagaacatg gtgacacgct atggcaagta cgaccccgaa ggtctcaact atgccaagat	1320
gatcctgtcc cacagggaag agctggacat caggacagtg tggaggacag aggagtacaa	1380
ggagaggacc agagtcttca acgaaatcac ggacttctcc agtgacctgc ccacgtcaaa	1440
ggcatggggc tggagagaca tagtcagagg aattcggaaa gtcgcagctc ctgtactgtc	1500
cacgtgtttt ccaatggcag caccactcat aggaatggca gaccaattca ttggagatct	1560
caccaagacc aacgcagcag gcggaaggta ccactccatg gccgcaggag ggcgctacaa	1620
agacgtgtc gagtccctgg caagcggagg gcccgacgga aaattctccc gagccctcaa	1680
gaacaggctg gagtccgcca actacgagga agtcgagctt ccacccccct caaaaggagt	1740
catcgtccct gtggtgcaca cagtcaagag cgcaccaggc gaggcattcg ggtccctggc	1800
aatcataatt ccaggggagt accccgagct tctagatgcc aaccagcagg tcctatccca	1860
cttcgcaaac gacaccggga gcgtgtgggg cataggagag gacataccct tcgagggaga	1920
caacatgtgc tacactgcac tccactcaa ggagatcaaa agaaacggga acatagtagt	1980
cgagaagatc tttgctggac caatcatggg tccctctgct caactaggac tgtccctact	2040
agtgaacgac atcaggagac gagttccaag gatggtattc accggcgaaa tcgccgatga	2100
cgaggagaca atcataccaa tctgcggtgt agacatcaaa gccatcgag cccatgaaca	2160
agggctgcc ctcatcggca accaaccagg agtggacgag gaggtgcgaa acacatccct	2220
ggccgcaçac ctgatccaga ccggaaccct gcccgtaaa cgcgcaaagg gctccaacaa	2280
gaggatcaag tacctgggag agctgatggc atcaaagca tccgggatgg acgaggaact	2340
gcaacgcctc ctgaacgcca caatggcacg ggccaaagaa gtccaggacg ccgagatcta	2400
caaacttctt aagctcatgg catggaccag aaagaacgac ctcaccgacc acatgtacga	2460
gtggtcaaaa gaggaccccg atgcactaaa gttcggaaaag ctcacagca cgccacaaa	2520
gcaccccgag aagcccaaag gaccagacca acaccatgcc caagaggcga gagccacccg	2580

```

catatcactg gacgccgtga gagccggggc ggacttcgcc acaccggaat gggtcgcgct 2640
gaacaactac cgcggcccat ctcccgggca gttcaagtac tacctgatca ctggacgaga 2700
accagaacca ggcgacgagt acgaggacta cataaaacaa cccattgtga aaccgaccga 2760
catgaacaaa atcagacgtc tagccaacag tgtgtacggc ctccacacc aggaaccagc 2820
accagaggag ttctacgatg cagttgcagc tgtattcgca cagaacggag gcagagggtcc 2880
cgaccaggac caaatgcaag acctcaggga gctcgcaaga cagatgaaac gacgaccccg 2940
gaacgccgat gcaccacgga gaaccagagc gccagcggaa ccggcaccgc ccggacgctc 3000
aaggttcacc ccagcggag acaacgctga ggtgtaacga ctactctctt tcctgactga 3060
tcccctggcc aaaaccccg ccccccaggg ggcccc 3097

```

```

<210> 3
<211> 972
<212> PRT
<213> Infectious pancreatic necrosis virus
<400> 3

```

```

Met Asn Thr Asn Lys Ala Thr Ala Thr Tyr Leu Lys Ser Ile Met Leu
1           5           10          15

```

```

Pro Glu Thr Gly Pro Ala Ser Ile Pro Asp Asp Ile Thr Glu Arg His
          20           25          30

```

```

Ile Leu Lys Gln Glu Thr Ser Ser Tyr Asn Leu Glu Val Ser Glu Ser
          35           40          45

```

```

Gly Ser Gly Ile Leu Val Cys Phe Pro Gly Ala Pro Gly Ser Arg Ile
          50           55          60

```

```

Gly Ala His Tyr Arg Trp Asn Ala Asn Gln Thr Gly Leu Glu Phe Asp
          65           70          75          80

```

```

Gln Trp Leu Glu Thr Ser Gln Asp Leu Lys Lys Ala Phe Asn Tyr Gly
          85           90          95

```

```

Arg Leu Ile Ser Arg Lys Tyr Asp Ile Gln Ser Ser Thr Leu Pro Ala
          100          105          110

```

```

Gly Leu Tyr Ala Leu Asn Gly Thr Leu Asn Ala Ala Thr Phe Glu Gly
          115          120          125

```

```

Ser Leu Ser Glu Val Glu Ser Leu Thr Tyr Asn Ser Leu Met Ser Leu
          130          135          140

```

```

Thr Thr Asn Pro Gln Asp Lys Val Asn Asn Gln Leu Val Thr Lys Gly
          145          150          155          160

```

```

Val Thr Val Leu Asn Leu Pro Thr Gly Phe Asp Lys Pro Tyr Val Arg
          165          170          175

```

Leu Glu Asp Glu Thr Pro Gln Gly Leu Gln Ser Met Asn Gly Ala Lys  
 180 185 190

Met Arg Cys Thr Ala Ala Thr Ala Pro Arg Arg Tyr Glu Ile Asp Leu  
 195 200 205

Pro Ser Gln Arg Leu Pro Pro Val Thr Ala Thr Gly Thr Leu Thr Thr  
 210 215 220

Leu Tyr Glu Gly Asn Ala Asp Ile Val Asn Ser Thr Thr Val Thr Gly  
 225 230 235 240

Asp Ile Asn Phe Ser Leu Thr Glu Gln Pro Ala Val Glu Thr Lys Phe  
 245 250 255

Asp Phe Gln Leu Asp Phe Met Gly Leu Asp Asn Asp Val Pro Val Val  
 260 265 270

Thr Val Val Ser Ser Val Leu Ala Thr Asn Asp Asn Tyr Arg Gly Val  
 275 280 285

Ser Ala Lys Met Thr Gln Ser Ile Pro Thr Glu Asn Ile Thr Lys Pro  
 290 295 300

Ile Thr Arg Val Lys Leu Ser Tyr Lys Ile Asn Gln Gln Thr Ala Ile  
 305 310 315 320

Gly Asn Val Ala Thr Leu Gly Thr Met Gly Pro Ala Ser Val Ser Phe  
 325 330 335

Ser Ser Gly Asn Gly Asn Val Pro Gly Val Leu Arg Pro Ile Thr Leu  
 340 345 350

Val Ala Tyr Glu Lys Met Thr Pro Leu Ser Ile Leu Thr Val Ala Gly  
 355 360 365

Val Ser Asn Tyr Glu Leu Ile Pro Asn Pro Glu Leu Leu Lys Asn Met  
 370 375 380

Val Thr Arg Tyr Gly Lys Tyr Asp Pro Glu Gly Leu Asn Tyr Ala Lys  
 385 390 395 400

Met Ile Leu Ser His Arg Glu Glu Leu Asp Ile Arg Thr Val Trp Arg  
 405 410 415

Thr Glu Glu Tyr Lys Glu Arg Thr Arg Val Phe Asn Glu Ile Thr Asp  
 420 425 430

Phe Ser Ser Asp Leu Pro Thr Ser Lys Ala Trp Gly Trp Arg Asp Ile  
 435 440 445

Val Arg Gly Ile Arg Lys Val Ala Ala Pro Val Leu Ser Thr Leu Phe  
 450 455 460  
 Pro Met Ala Ala Pro Leu Ile Gly Met Ala Asp Gln Phe Ile Gly Asp  
 465 470 475 480  
 Leu Thr Lys Thr Asn Ala Ala Gly Gly Arg Tyr His Ser Met Ala Ala  
 485 490 495  
 Gly Gly Arg Tyr Lys Asp Val Leu Glu Ser Trp Ala Ser Gly Gly Pro  
 500 505 510  
 Asp Gly Lys Phe Ser Arg Ala Leu Lys Asn Arg Leu Glu Ser Ala Asn  
 515 520 525  
 Tyr Glu Glu Val Glu Leu Pro Pro Pro Ser Lys Gly Val Ile Val Pro  
 530 535 540  
 Val Val His Thr Val Lys Ser Ala Pro Gly Glu Ala Phe Gly Ser Leu  
 545 550 555 560  
 Ala Ile Ile Ile Pro Gly Glu Tyr Pro Glu Leu Leu Asp Ala Asn Gln  
 565 570 575  
 Gln Val Leu Ser His Phe Ala Asn Asp Thr Gly Ser Val Trp Gly Ile  
 580 585 590  
 Gly Glu Asp Ile Pro Phe Glu Gly Asp Asn Met Cys Tyr Thr Ala Leu  
 595 600 605  
 Pro Leu Lys Glu Ile Lys Arg Asn Gly Asn Ile Val Val Glu Lys Ile  
 610 615 620  
 Phe Ala Gly Pro Ile Met Gly Pro Ser Ala Gln Leu Gly Leu Ser Leu  
 625 630 635 640  
 Leu Val Asn Asp Ile Glu Asp Gly Val Pro Arg Met Val Phe Thr Gly  
 645 650 655  
 Glu Ile Ala Asp Asp Glu Glu Thr Ile Ile Pro Ile Cys Gly Val Asp  
 660 665 670  
 Ile Lys Ala Ile Ala Ala His Glu Gln Gly Leu Pro Leu Ile Gly Asn  
 675 680 685  
 Gln Pro Gly Val Asp Glu Glu Val Arg Asn Thr Ser Leu Ala Ala His  
 690 695 700  
 Leu Ile Gln Thr Gly Thr Leu Pro Val Gln Arg Ala Lys Gly Ser Asn  
 705 710 715 720  
 Lys Arg Ile Lys Tyr Leu Gly Glu Leu Met Ala Ser Asn Ala Ser Gly

					725					730								735
Met	Asp	Glu	Glu	Leu	Gln	Arg	Leu	Leu	Asn	Ala	Thr	Met	Ala	Arg	Ala			
			740						745				750					
Lys	Glu	Val	Gln	Asp	Ala	Glu	Ile	Tyr	Lys	Leu	Leu	Lys	Leu	Met	Ala			
		755					760					765						
Trp	Thr	Arg	Lys	Asn	Asp	Leu	Thr	Asp	His	Met	Tyr	Glu	Trp	Ser	Lys			
	770					775					780							
Glu	Asp	Pro	Asp	Ala	Leu	Lys	Phe	Gly	Lys	Leu	Ile	Ser	Thr	Pro	Pro			
785					790					795					800			
Lys	His	Pro	Glu	Lys	Pro	Lys	Gly	Pro	Asp	Gln	His	His	Ala	Gln	Glu			
				805					810					815				
Ala	Arg	Ala	Thr	Arg	Ile	Ser	Leu	Asp	Ala	Val	Arg	Ala	Gly	Ala	Asp			
			820					825					830					
Phe	Ala	Thr	Pro	Glu	Trp	Val	Ala	Leu	Asn	Asn	Tyr	Arg	Gly	Pro	Ser			
		835				840						845						
Pro	Gly	Gln	Phe	Lys	Tyr	Tyr	Leu	Ile	Thr	Gly	Arg	Glu	Pro	Glu	Pro			
	850					855					860							
Gly	Asp	Glu	Tyr	Glu	Asp	Tyr	Ile	Lys	Gln	Pro	Ile	Val	Lys	Pro	Thr			
865					870					875					880			
Asp	Met	Asn	Lys	Ile	Arg	Arg	Leu	Ala	Asn	Ser	Val	Tyr	Gly	Leu	Pro			
				885					890					895				
His	Gln	Glu	Pro	Ala	Pro	Glu	Glu	Phe	Tyr	Asp	Ala	Val	Ala	Ala	Val			
			900					905					910					
Phe	Ala	Gln	Asn	Gly	Gly	Arg	Gly	Pro	Asp	Gln	Asp	Gln	Met	Gln	Asp			
		915					920					925						
Leu	Arg	Glu	Leu	Ala	Arg	Gln	Met	Lys	Arg	Arg	Pro	Arg	Asn	Ala	Asp			
	930					935					940							
Ala	Pro	Arg	Arg	Thr	Arg	Ala	Pro	Ala	Glu	Pro	Ala	Pro	Pro	Gly	Arg			
945					950					955					960			
Ser	Arg	Phe	Thr	Pro	Ser	Gly	Asp	Asn	Ala	Glu	Val							
				965					970									

```
<210> 4
<211> 26
<212> DNA
<213> Artificial sequence
<220>
```

<223> Synthetic construct

<400> 4

aactcgagat ggtgagcaag ggcgag

26

<210> 5

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 5

atctcgactt gtacagctcg tccatgc

27

<210> 6

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 6

gggatccatg aacacaaaca aggc

24

<210> 7

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 7

aaagcttaca cctcagcggt gtc

23

<210> 8

<211> 5170

<212> DNA

<213> Aequorea victoria

<220>

<221> misc feature

<222> (651)..(651)

<223> n can be a, g, t or c

<400> 8

aagcttcaaa ttaagtcagc tccttaaagt aaagataata aagtgtagtt caagaactat 60

atgaatgatg tgttttcaga taacccaaat ggggaaaaac atgctaaagt cagcatatatt 120

ttggaaaatt gatgacgtca tcatgacgtc gttttgatga caaaacttat tataagcgaa 180

ttcttatatt ttacaggat aacaaagatg agtaaaggag aagaactttt cactggagtt 240

gtcccaattc ttgttgaatt agatgggtgat gtaaatgggc acaaatcttc tgtcagtggg 300

gagggtgaag gtgatgcaac atacggaaaa cttaccctta aatttatttg cactactgga 360

aagctacctg ttccatggcc aacacttgct actactttct cttatgggtg tcagtaagtg 420

cattttatac tcttttaata tcagtgttaa gaaaatcaag tgtcttgcta ttttttcgat 480



tattggtgca attctagtca aattattgcg tttttttacc caaatgtta atgtaaaact	540
gaaatttggc acacttgccg aaatatatac agggatattt gaaaaatta aacaggatga	600
taaaagttgc acagaaactt atctcaagat ttacccgcag aaagatgctt naaaaattga	660
tatttgacag agcaaaacct gagattcacg tcttttagtt gtttgacttg aaattttggt	720
gacaggtagg tatcatgaaa aacaaacaaa acgtaaaaat atcacgtgat taaagtgtat	780
cttacagacc agaaacagtt ttattaactt ctattattct attttgcaat atacacattg	840
tatcaatttc ttgagttact cgaagtaata ccgacctatc atcagaattt caagtcaaca	900
caacattata tggggctgat tagggaatga ttttgtctct tttagatgct tttcaagata	960
cccagatcat atgaaacagc atgacttttt caagagtgcc atgccgaag gttatgtaca	1020
ggaaagaact atattttaca aagatgacgg gaactacaaa tcacgtgctg aagtcaagtt	1080
tgaaggatgat accctcgta atagaattga gttaaaagggt attgatttta aagaagatgg	1140
aaacatttct ggacacaaaa tggaatacaa ctataactca cacaatgtat acatcatggc	1200
agacaaacaa aagaatggaa tcaaagttaa cttcaaaatt gtatgtatac gttaagggca	1260
taaatttttg cgggcataaa atcttgcgaa atttattatc gcgaatagggt tacgcaaat	1320
ctataattaa aatgtatttt tttctgctga ttttctaaat aacaactcaa cccgtcattt	1380
ttatatcgca aaaataaatt ccgaaataat ttatgctcgc aaasatttag gcccataagt	1440
agacttttga tatctgctg ctctgcaatg aagtaaaaat acgatatttt cattgaaata	1500
cacgggttca aagttatttg ttaattcaat aagcgtgcgc agaaattaaa ggacgtataa	1560
agatacgaac acatcaaacc attcatgcgt aaataatgtt ctatttttaa aattcaccaa	1620
agcttaaata ttcttaagaa ttattcatgt gccatgggag caacaatata gttatggaca	1680
aaaatttctg agttcacttt tatttctgcg cgcccgcatc aaagttcaaa caactgtgaa	1740
cccagatttt ttccagcttg caattttaat aagagacaaa aagcaaattg cagttcaaga	1800
aaatcgagat attgccagat gtaaacattt aataagagac aaaaagttca taagcgttct	1860
aaagaacagc aacaaaataa taattagaat taaacgagtt ctcaaacaaa ataaaaactg	1920
aagtcaaaga gtcagtaagg aatttagtta acgatgcttt ataatacaag ttttaattcc	1980
agttcatgta tgcaattaac aataagatct tggagaattg aatatgtttc gaaattttat	2040
aaattcggat ttaatttcta aagttgtgta tcaaaaatag ttcaaaactat tttcatgaaa	2100
agatgataaa ttacggtaat aagtatataa tataatcaat taaaattaat tttaggtca	2160
aattacagaa tccacgtttt ttttctctag acatagcaca gtgttttagat gtttgtttta	2220
tttcatccat ccttattaca gttttctct gaactttaat actagcgtac aatttgaata	2280
ataatctgaa atgattcaac ttttcagaga cacaacattg aagatggaag cgttcaacta	2340
gcagaccatt atcaacaaaa tactccaatt ggcgatggcc ctgtcctttt accagacaac	2400
cattacctgt ccacacaatc tgccctttcc aaagatccca acgaaaagag agatcacatg	2460
atccttcttg agtttgaac agctgctggg attacacatg gcatggatga actatacaaa	2520

taaatgtcca gacttccaat tgacactaaa gtgtccgaac aattactaaa atctcaggg	2580
tcctgggttaa attcaggctg agataattatt tatatatatta tagattcatt aaaattttat	2640
gaataattta ttgatgttat taataggggt tttttctta ttaaataaggc tactggagtg	2700
cattcctaatt tctatattaa ttacaatttg atttgacttg ctcagaatcc cgcttcattg	2760
cttttccact tgcattatcc ttatttagta ttaatttgta ttttggttg gctacattga	2820
gtgcaaaaaa cctaattttc ggacgaattt tcgaacgaat ttttttgacg gaattttctt	2880
cattctatatt actcctctag cttaaattatt ttaccttttt gttaatttg ttaaattatt	2940
ctctgagccg atgattgaga aattaatgga ttaaaagtga gtacctaca tgttgtcaac	3000
ttgtaacgaa tggaaaaaga aattacgttt caagagtttg aaaggtaata cagttacagt	3060
taaccgcaga aaaattgcat gatgattgat aaattcgatt tttgttatcc taaaattttc	3120
caaacgtcag tggccgacga ctttatcagg gacttctaaa agtgaaaaat aatcagggtgc	3180
ggatttcgaa ggcgcaaac tataggaaga gagcgaaatg tcattaaatt atcatattct	3240
attaactgat gacaatagat gatgaaaagt ttatgattat tcaactctct cctgtaatta	3300
tgcgaccott ctagattcac gcctgaaagt atagctacct gggatgaagt actagtctga	3360
ggactcttca cctaaaaatt aaattcttat aagagtaaac aagaaactta gcagttacaa	3420
acgggagagc gatgagaaac aaaaacaatt acgttgccac tatgaatatc gatgttcaat	3480
caattttgtt cttacttat aagaacgaga tcgtcttaac ttaaaatagt aaaatgttat	3540
caagataata gcaatttttt accgacacag cgaagactca ctactgaaat gatcagtttt	3600
aatcaggcaa ataatocgtg gcacataata gtgaccgaaa ataattaatc ggcattaaga	3660
ctaccgaaat aataatgttt tttctactgc gtatacgcgt gagaaatttt caataagctc	3720
atcatcttca gcatagtatt acttttatgt aaagtatcaa ttccgacata aaataacggc	3780
ttattatcga aataatagcg ttttctctac tccatgcgcg tcaaaagtgc tctctaggct	3840
catcatcttc agcataatta taatttttgt aaagtaccag ttccggtcga aaataatgac	3900
taattaccga aattatagtg tttttctatt gccatgcgcg tgaaaaattt tgattgaatc	3960
atcatcttca gcataggcat aattctttgt aaaatatcga ttccgacata aaataatggc	4020
ctattaccga aataatcgcg tttttctctac tgcgcatgcg cgtcaaaaat tatattttta	4080
ttcatcatct tcagcataat tatatttttt tgtaaagtac cagttccggt agaaaataat	4140
gacttggtac tgaaataata gcgtttttct attgcgcgtg cgctataaaa attaaagtaa	4200
cgtcatcata ttcagcatgg tattgaaatt ttcaaattta attaacctat tgaacaagaa	4260
tgtacacttg catcaaaata ggtgaaattc gccaatatcg ctaaatgtga cgcgcgagg	4320
caatactacg catgtagctt caggtaaagc atgtagaaac tcggaggagt aggagtccac	4380
cgtcgaaaact aaaacgggat acactacgct atggccttcg ctctcccgta aaaagggact	4440
aacaatacga cctaattgaa atactaaaaa aaacaagaga aatttaaccc ctttggttaac	4500
acttttcaaa agtgggattt tttagccaac catctgggtat atatggttgc tcattttatt	4560
attatctctt tctttattgt tggatcaacg tagtcaaaat acaaattagg ttaataaaaa	4620

gcaacattat aatgtataaa atctaattgt gtctaattac cgacaaattt tacaggaaca 4680  
 gttttcacca gaccgagtct taatttttagt tttaaaagaa attatgtttc tactgtttctg 4740  
 acaatctgaa gacaattagt tctagtgtaa caatgctctg aattgaatat attcagcaat 4800  
 attttgtttg taagaattgg atgaatgtac gaaccttcag cagattttata ccaagtgtta 4860  
 gatttaacaa gatttgcaag ctgatgagtt tcgagaaaaat tcaacatatac tggatttgag 4920  
 ggtggaacat taaaatctcc taagataata attctatcat aattagaata taaattatca 4980  
 atgatgtcat ttaagtgtac tagaaaaata ttgatagtaa cagttggatg tttgtatata 5040  
 gaaatagtaa gccatctatt tttcccaaat gcgagttcaa aaacccaaat tggattcctt 5100  
 caaagaaaaa agacattaag aaacttgatg gaatcccttc tcgactgtaa acaagcagtc 5160  
 tctgggatcc 5170

<210> 9  
 <211> 966  
 <212> DNA  
 <213> Aequorea victoria

<400> 9  
 tacacacgaa taaaagataa caaagatgag taaaggagaa gaacttttca ctggagtgtg 60  
 cccaattctt gttgaattag atggtgatgt taatgggcac aaattttctg tcagtggaga 120  
 ggggtgaaggt gatgcaacat acggaaaact tacccttaaa tttatttgca ctactggaaa 180  
 actacctgtt ccatggccaa cacttgtcac tactttctct tatggtgttc aatgcttttc 240  
 aagataccca gatcatatga aacagcatga ctttttcaag agtgccatgc ccgaaggtta 300  
 tgtacaggaa agaactatat ttttcaaaga tgacgggaac tacaagacac gtgctgaagt 360  
 caagtttgaa ggtgataccc ttgttaatag aatcgagtta aaagggtattg attttaaaga 420  
 agatggaaac attcttggac acaaattgga atacaactat aactcacaca atgtatacat 480  
 catggcagac aaacaaaaga atggaatcaa agttaacttc aaaattagac acaacattga 540  
 agatggaagc gttcaactag cagaccatta tcaacaaaat actccaattg gcgatggccc 600  
 tgtcctttta ccagacaacc attacctgtc cacacaatct gccctttcga aagatcccaa 660  
 cgaaaagaga gaccacatgg tccttcttga gtttgtaaca gctgctggga ttacacatgg 720  
 catggatgaa ctatacaaat aaatgtccag acttccaatt gacactaaag tgtccgaaca 780  
 attactaaaa tctcagggtt cctgggttaa ttcaggctga gatattattt atatatttat 840  
 agattcatta aaattgtatg aataatttat tgatgttatt gatagaggtt attttcttat 900  
 taaacaggct acttggagtg tattcttaat tctatattaa ttacaatttg atttgacttg 960  
 ctcaaa 966

<210> 10  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 10  
taatacgact cactatagga aagagagttt caacg 35

<210> 11  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Synthetic Construct

<400> 11  
ggccatggag tggtagcttc 20

<210> 12  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Synthetic Construct

<400> 12  
gaaggtacca ctccatggcc 20

<210> 13  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Synthetic Construct

<400> 13  
aaagcttctg cagggggccc cctggggggc 30

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/25185

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :A61K 39/12, 39/295; C12N 5/00, 15/09, 15/00, 15/74, 7/00, 7/04

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/199.1, 202.1, 204.1, 205.1; 435/235.1, 69.3, 236, 472, 320.1, 325

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	US 6,274,147 B1 (VAKHARIA et al.) 14 August 2001, please see entire document	1-24
Y	US 6,180,614 B1 (DAVIS) 30 January 2001, please see entire document.	1-24
Y	US 5,939,073 A (MCLOUGHLIN et al.) 17 August 1999, see entire document.	1-24



Further documents are listed in the continuation of Box C.



See patent family annex.

"	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

23 SEPTEMBER 2002

Date of mailing of the international search report

11 DEC 2002

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

LAURIE SCHEINER

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/25185

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/199.1, 202.1, 204.1, 205.1; 435/235.1, 69.3, 236, 472, 320.1, 325